

47

Table of Contents

(NOTE: This document is incomplete. 100 pages, including all chapters beyond 0 5 a) are missing, as are illustrations beyond No. 30, and the entire appendix.)

I. Definition of the region

II. Geological summary

A. Upper Rhenish plains

B. Old mountain sections on both sides of the Rhine Valley Trough

1. Northern structural elements

- a) Saar - Nahe region
- b) Palatinate Forest
- c) Odenwald
- d) Spessart

2. Southern structural elements

- a) Vosges
- b) Black Forest

C. Recent covering mountains

1. Lorraine steps

2. Swabian - Franconian steps

III. Ore deposits

A. Rhine Valley Trough

1. Tertiary contact-pneumolytical niobium deposit in the
Kaiserstuhl

2. Tertiary ore deposits

- a) Eocene deposits in Kandern District
- b) Pliocene deposits in the Mainz basin

3. Quarternary gold washings

B. Saar - Nahe region

1. Epithermal copper ore deposits

2. Epithermal mercury deposits

- a) Lemberg mine in the Niederhaeuser Forest on the Nahe River

48

b) Landsberg near Ober-Noschel

c) Stahlberg near Katzenbach

3. Epithermal barite deposits

4. Telethermal copper ores

5. Sedimentary iron ore deposits

a) Upper Carboniferous deposits

b) Lower Rotliegendes deposits

c) Limonite ores in Red Triassic sandstone

C. Odenwald and Spessart

1. Epithermal beld-copper-pyrite deposits along the Bergstrasse

2. Telethermal metasomatic zinc-lead ore deposits at Wiesloch

3. Epithermal copper deposits of Reichenbach

4. Epithermal to telethermal silver-cobalt-nickel deposits

5. Epithermal specularite-hematite veins

6. Epithermal to telethermal barite deposits

a) Barite veins in crystalline rock

b) Barite veins in Rotliegendes

c) Barite veins in Permian limestone

d) Barite veins in Red Triassic sandstone

7. Iron ores in Rotliegendes

8. Metasomatic manganese-iron and iron ores in Permian limestone

9. Copper, lead, and silver ores in copper clay of Permian limestone

D. Vosges

1. Pegmatitic-pneumatolytic molybdenum deposits

2. Pneumatolytic-catathermal iron ore deposits

a) Tourmaline-bearing specularite-magnetite veins at Rothau

b) Specularite-magnetite veins at Framont

3. Mesothermal arsenic ore veins

4. Mesothermal to epithermal lead-zinc-silver ore veins

a) Carboniferous spathic lead-zinc-silver veins of Markirch

- b) Carboniferous spathic lead-zinc-silver veins of St. Kreuz
 - c) Lead-zinc-silver veins in the upper Weiler valley
 - d) Lead-zinc-silver veins in the upper Masmuenster valley
 - e) Lead-zinc-silver veins of Giromagny
 - 5. Epithermal copper ore veins, to some extent with cobalt-nickel ores
 - a) Tetrahedrite-chalcopyrite veins of Markirch
 - b) Tetrahedrite-chalcopyrite veins in the upper Weiler valley
 - c) Chalcopyrite veins in the upper Thanner valley
 - d) Chalcopyrite veins of Plancher-les-Mines
 - 6. Epithermal antimony ore veins
 - 7. Epithermal iron and manganese ore veins
 - a) Siderite veins converted into limonite veins
 - b) Specularite veins
 - 8. Epithermal fluorite veins
 - 9. Epithermal barite veins
- E. Black Forest
- 1. Liquid magmatic nickel deposits
 - 2. Pneumatolytic cassiterite deposits
 - 3. Hydrothermal ore veins
 - a) Mesothermal to epithermal lead-zinc-copper-arsenic ore veins in the Muenster valley
 - (1) Ore veins north of Belchen Mountain
 - (2) Ore veins north and south of the Muenster valley
 - (3) Ore veins on the western slope of the Muenster valley
 - b) Mesothermal fluorbaritic copper-bismuth veins and epithermal cobalt-nickel-bismuth-silver veins at Wittichen
 - c) Mesothermal lead-copper-silver veins at Oberwolfach
 - d) Mesothermal to epithermal lead-silver veins in the Schapbach region

- e) Mesothermal silver-quartz veins at Bollenbach-Haslach
- f) Epithermal lead-zinc veins on Schauinsland Mountain
- g) Epithermal to telethermal lead-zinc veins at Badenweiler
- h) Epithermal lead-zinc veins between Belchen Mountain and Todtnau
- i) Epithermal lead ore veins near St. Blasien and Birkendorf
- k) Epithermal lead-zinc ore veins in the Dreisam and Kandel regions
- l) Epithermal to telethermal lead-zinc vein east of Lehr
- m) Epithermal lead-zinc veins in the lower Kinzig valley
- n) Epithermal lead-zinc veins at Hausach
- o) Epithermal baritic lead-copper veins at Rippoldsau
- p) Epithermal iron-manganese ore veins at Eisenbach and Voehrenbach
- q) Epithermal iron ore veins in the Kandel region
- r) Epithermal iron and manganese ore veins in the lower Kinzig region
- s) Epithermal limonite veins near Oberwolfach
- t) Epithermal fluorite veins
- u) Epithermal barite veins
- v) Telethermal copper, bismuth, and barite veins at Freudenstadt
- w) Telethermal copper-bismuth veins at Neubulach
- x) Telethermal barite-iron ore and fluorite veins of Nauenburg-Pforzheim
- y) Telethermal barite veins

F. Lorraine steps

1. Telethermal iron ore veins
2. Sedimentary lead ore veins
3. Sedimentary iron ore veins
 - a) Lorraine-Luxemburg minette ores
 - b) Various minor iron ore deposits

51

G. Swabian-Franconian steps

1. Epithermal fluorite veins
2. Telethermal barite veins
3. Lead ore deposits in the Keuper
4. Oolithic iron ores in the Jurassic
 - a) Lias ores
 - b) Dogger ores
- 5) Iron ores in the Cretaceous and in the Tertiary
 - a) Pre-Cenomanian Cretaceous ores in the "Franconian Alb" Mountains
 - b) Dogger ores in the Franconian Upper Cretaceous - Tertiary, reassorted several times
 - c) Tertiary and diluvial bean ores

IV. The geological and structural factors in their relationships to the formation of deposits

A. Spatial distribution of the deposits in the individual areas

1. Rhine Valley Trough
2. Saar - Nahe region
3. Odenwald - Spessart
4. Vosges
 - a) Northwestern and eastern slopes of the Champ du Feu massif (Hochfeld)
 - b) Markirch-Urbeis region
 - c) Region of the Alsatian Belchen Mountains
5. Black Forest
 - a) Murgtal granite region
 - b) Triberg granite region
 - c) Eisenbach granite region
 - d) Zone of denuded granite anticlines in the southern Black Forest
6. Lorraine steps
7. Swabian - Franconian steps

B. Spatial arrangement of the deposits in the entire region

C. Temporal arrangement of the formation of the deposits

V. Summary

Bibliography

Illustrations in the Text

Fig. 1: Tectonic sketch of the Black Forest, the Vosges, and the Rhine Valley Trough

Fig. 2: Schematic sketch of the edge of the Black Forest at Freiburg in Breisgau

Fig. 3: Synclines and rises of the Varistian orogenesis

Fig. 4: Cross-section of the Queich valley near Albersweiler in the Palatinate

Fig. 5: Structural map of the Odenwald

Fig. 6: Cross-section of the Spessart foothills

Fig. 7: Tectonic cross-section of the overthrust structure of the Vosges

Fig. 8: Tectonic sketch of the Vosges and the Westrich syncline

Fig. 9: Tectonic map of the Black Forest

Fig. 10: Bedding of the granites and Carboniferous sediments in the southern Black Forest

Fig. 11: Sections of the Keuper and Jurassic stages in Swabia and Franconia

Fig. 12: Sketch of the bedding of the rocks in Swabia and Franconia.
Contour lines of the border between the Franconian deposits and the lettenkohle

Fig. 13: Distribution of the bean ore deposits in Rhenish Hesse

Fig. 14: Copper ore deposits in the Saar

Fig. 15: Iron ore deposits in the Saar

Fig. 16: Map of the ore region near Wiesloch

Fig. 17: Sketch of ore deposit IV in the Kobelsberg near Wiesloch, with two vertical sections

Fig. 18: Sketch of the southern part of ore deposit V in the Kobelsberg, near Wiesloch

Fig. 19: The barite veins around Grosse Auersberg Mountain, northeast of Brusckensau

Fig. 20: Facies and thickness conditions of the Dogger in South Germany

Fig. 21: Facies of the oolith bank in the Lower Lias of Swabia

Fig. 22: Facies of the copper rock bank in the Lower Lias of Swabia

Fig. 23: Distribution of Lias ores on the western edge of the Bohemian Mass

Fig. 24: Facies conditions in the Dogger in Alsace, Baden, and Swabia

Fig. 25: Paleographic map of the Dogger sandstone and the iron ooliths in the syncline of the Franconian Alb Mountains

Fig. 26: Facies conditions in the Dogger in Baden and Swabia

Fig. 27: Paleographic map of the subfurcate and Humphries strata in the syncline of the Franconian Alb Mountains

Fig. 28: Sketch showing the principle of the formation of iron ooides

Fig. 29: Cross-section through the ore mountain of Amberg. Basal and overthrust parts of the ore body

Fig. 30: The bean and Dogger ores from the center of the anticline of Central Franconia near Eichstaett and Neumarkt

Fig. 31: Schematic representation of the formation of iron ore deposits in the Nitzelbuch mine near Auerbach

Fig. 32: The daily output of mineral water by the mineral springs of Germany

Fig. 33: The concentration of free CO in the mineral springs of Germany

Fig. 34: The daily output of CO by the mineral springs of Germany

Fig. 35: The North Rhenish constriction of the Rhenish shield

Fig. 36: Hypothetical and schematic representation of the contour lines of the Rhenish shield

51

Illustrations in Appendix

- Fig. 1: Definition of the region. The Rhine Valley Trough with its border regions and the adjacent step topography
- Fig. 2: Map of the Rhenish shield with the Rhine Valley Trough and the volcanic belts
- Fig. 3: Geological map of the Naha syncline and the Rhenish Palatinate with copper and mercury deposits
- Fig. 4: Map of deposits and structure of the Odenwald
- Fig. 5: Geological and deposit map of the Spessart
- Fig. 6: Geological and deposit map of the Vosges
- Fig. 7: Geological map of the Central Black Forest
- Fig. 8: Map of the granite mass of the southern Black Forest
- Fig. 9: Map of the mineralization zone of the northern Black Forest
- Fig. 10: Map of the mineralization zone of the southern Black Forest
- Fig. 11: Geological and deposit map of the Lorraine steps
- Fig. 12: Geological and deposit map of the Swabian-Franconian steps
- Fig. 13: The sedimentary syngenetic iron ore deposits and mineral pigments of the northern Franconian Alb Mountains
- Fig. 14: Geological map of the Kaiserstuhl Mountains
- Fig. 15: The copyrite deposits near Scheelingen
- Fig. 16: Map showing the relationships between the mobilized telethermal deposits and the volcanic and carbon dioxide belts of the Rhenish shield
- Fig. 17: Map of the gravity deviations of South Germany, according to K. Schuette
- Fig. 18: Table of the temporal sequence of the deposit formation